

Magnesium and calcium excretion during pregnancy

By L. Spätling, P.A. Kunz*, R. Huch, A. Huch

Departement of Obstetrics, University Hospital, Director: Prof. Dr. A. Huch, University of Zürich, Switzerland

*Medical Chemical Central Laboratory of the University Hospital, Director: Prof. Dr. D. J. Vonderschmitt, University of Zürich, Switzerland

Zusammenfassung

Zur Untersuchung der Gründe des Absinkens des Magnesiumplasmagehaltes (Mg) in der Schwangerschaft, von dem gezeigt werden konnte, daß er mit einer Erniedrigung des Mg-Spiegels im Myometrium korreliert, sollte die Mg- und Kalziumausscheidung (Ca) im Urin in der Schwangerschaft bestimmt werden. Es wurde bei sieben Probandinnen während der Schwangerschaft in vierzehntägigen Abständen 24-Stunden-Urin gesammelt und Blut entnommen. Im Urin und Plasma wurde Mg mit Atomabsorptionsspektrophotometrie und Ca colorimetrisch bestimmt. Die Clearances wurden errechnet. Es zeigte sich eine 25%ige Erhöhung der Mg-Ausscheidung auf 4,78 mmol/24 h ($p < 0,05$). Der Mg-Plasmawert sank um 25 % auf 0,70 mmol/l ($p < 0,01$). Die Mg-Clearance wurde um 41 % auf 4,57 ml/min erhöht ($p < 0,05$). Schon im ersten Trimenon ist die Ca-Ausscheidung um 41 % auf 4,86 mmol/24 h angestiegen ($p < 0,05$). Die Ca-Plasmakonzentration fällt um 12 % auf 2,16 mmol/l ($p < 0,001$). Die Ca-Clearance steigt um 123 % auf 2,03 ml/min an ($p < 0,02$). Somit ist wahrscheinlich die Erhöhung der Mg-Ausscheidung im Urin für eine Mg-Mangelsituation mitverantwortlich. Für die relativ geringere Rückresorption kommen schwangerschaftsspezifische Veränderungen in Frage: 1. Eine Erhöhung der extrazellulären Flüssigkeit, 2. die Erhöhung der glomerulären Filtrationsrate (GFR) und 3. die durch die vermehrte Natrium-(Na)-Reabsorption reduzierte Mg- und Ca-Reabsorption.

Summary

The magnesium (Mg) and calcium (Ca) excretion in the urine during pregnancy was determined in order to examine the reason for the decrease in the magnesium plasma level during pregnancy, a decrease which was shown to correlate with the diminished Mg-level in the myometrium. 24-hour urine and blood samples were taken from seven pregnant volunteers once every two weeks. The Mg was determined in the urine and plasma by using atom absorption spectrometry and the Ca was determined by colorimetry. The clearances were calculated. A 25 % increase in Mg-excretion to 4.78 mmol/24 h ($p < 0.05$) was shown. The Mg-plasma level diminished by 25 % to 0.70 mmol/l ($p < 0.01$). The Mg-clearance increased by 41 % to 4.57 ml/min ($p < 0.05$). Already in the first trimester

the Ca-excretion increased by 41 % to 4.86 mmol/24 h ($p < 0.05$). The Ca-plasma concentration diminished by 12 % to 2.16 mmol/l ($p < 0.001$). The Ca-clearance increased by 123 % to 2.03 ml/min ($p < 0.02$). The increased Mg-excretion in the urine is therefore probably one of the reasons for Mg-deficiency. The following changes associated with pregnancy are to be considered as reasons for the relatively low reabsorption: 1. The increase in extracellular fluid, 2. the increase in glomerular filtration rate (GFR), and 3. reduced Mg and Ca-reabsorption due to increased sodium (Na) reabsorption.

Résumé

Les excréations de magnésium (Mg) et de calcium (Ca) urinaires ont été étudiées au cours de la grossesse dans le but de rechercher les raisons de la baisse gravidique du Mg plasmatique, réduction corrélée avec la baisse du Mg myométrial. Des prélèvements des urines de 24 h et du sang ont été effectués chez 7 femmes enceintes volontaires, chaque 2 semaines. Le Mg a été déterminé par spectrométrie d'absorption atomique et le Ca par colorimétrie. Les clairances ont été calculées. Un accroissement de 25 % de la magnésurie à 4,78 mmol 24 h ($p < 0,05$) a été observé. Le taux de Mg plasmatique est réduit de 25 % à 0,70 mmol/l ($p < 0,01$). La clairance du Mg augmente de 41 % à 4,57 ml/min ($p < 0,05$). L'excrétion calcique augmente de 41 % pendant le 1^{er} trimestre à 4,86 mmol/24 h ($p < 0,05$). La concentration de Ca plasmatique diminue de 12 % à 2,16 mmol/l ($p < 0,001$). La clairance du calcium augmente de 123 % à 2,03 ml/min ($p < 0,02$). L'augmentation de la magnésurie est donc probablement une des raisons de la déficience magnésique. Elle peut provenir: 1. d'une augmentation des liquides extra-cellulaires, 2. d'une augmentation de la filtration glomérulaire, et enfin, 3. d'une réabsorption réduite de Ca et Mg due à une réabsorption accrue du Na.

Introduction

The observation that premature contractions can be reduced partly by administering Mg (Spätling 1981), drew attention to the role of Mg in pregnancy. As more precise methods of analysis such as atom absorption spectrometry became more wide-spread, it became feasible to correlate the data from these analyses to clinical manifestations. Baltzer and Daume, 1976, and others (De Jorge et al., 1965, and Brockerhoff et al., 1981) observed a reduction in Mg-plasma level particularly at the beginning of pregnancy. The Mg-plasma level reduced consistently and over a long period of time seems to influence growing tissues because it could be shown that the Mg-concentration of the uterus musculature diminished during the course of pregnancy (Spätling et al., 1983). Although the Mg-requirement of the uterus and foetus is not insignificant, it is unlikely that consistently low Mg-plasma values can be explained simply in these terms. The hypothesis established for the present study therefore was that urinary Mg-excretion increases during pregnancy. For this purpose, urinary Mg-excretion, Mg-clearance and Mg-plasma levels were measured during normal pregnancy and compared to the corresponding Ca-values.

Materials and methods

Seven volunteers aged 17 to 36 ($n = 26$) participated in the study. Four of the women had their first child, two their second and one her third. They joined the study between their sixth and one her third. They joined the study between their sixth and twelfth

Na remains constant while potassium and chloride rise.

The changes in protein binding of Mg and Ca in pregnancy are difficult to interpret in this connection. If it is calculated only with the decrease in plasma protein concentration, a decrease of ionized Mg of 10% and of Ca of 7% is found none the less. It is unlikely that pregnancy does not affect the fraction of ionized Mg and Ca.

Heaney and Skillman (1971) observed a Ca-excretion of 3.49 mmol/24 h in non-pregnant women, similar to the 6–8 week postpartal value we found. Davis et al. (1970) found a slightly higher level of excretion at 4.55 mmol/24 h. Increased Ca-excretion in pregnancy with the exception of the last five weeks is described by Heaney and Skillman (1971). We are not able to confirm this finding in our study.

One reason for the increased Mg and Ca-excretion during pregnancy may be the 30% increase in glomerular filtration rate (GFR) (Sim and Krantz, 1958). Massry (1981) writes that a constant increase of GFR could cause a significant decrease of Mg-plasma level. Another reason for the increased excretion of Mg and Ca could be the increase in extracellular fluid caused by pregnancy. Massry et al. (1969) describe a decrease in the tubular reabsorptive capacity caused by an increase of extracellular volume. The studies of Di Bona et al. (1971), demonstrating reduced Na-reabsorption after MgSO₄ infusion, clearly show how the reabsorptions of Mg, Ca and Na influences each other in the distal portion of the upward Henle loop. Massry et al. (1970) consider that the sites of reabsorption are limited and they demonstrate increased Mg, Na and Ca excretion in the urine after a MgCl₂ infusion. The increase in GFR due to pregnancy necessitates an additional Na-reabsorption of 10 000 mmol/24 h. It is therefore conceivable that this

significant increase in Na-reabsorption occurs at the expense of Mg- and Ca-reabsorption, of which only 50 mmol/24 h, respectively 160 mmol/24 h, must be reabsorbed additionally during pregnancy.

The 3 mmol/24 h increase in Ca excretion that we measured corresponds to an approx. 10 mmol/24 h increase of Ca-reabsorption during pregnancy. No similar studies have been conducted on the balance of Mg, but the 25% increase in Mg-excretion we measured seems to manifest itself in the cell Mg content of pregnant women. As mentioned before, we were able to show a decrease in the Mg-level of myometrial cells during the course of pregnancy (Spätling et al., 1983), but not a change in the Ca-level (Spätling et al., to be published).

The role of the parathyroid hormone, which is important in regulating Ca and Mg overall, is unclear because the level of this hormone increases only after the 24th to the 28th week (Cushard et al., 1972). The decrease in the Mg and Ca level, however, can be observed much earlier (Brockhoff et al., 1981).

The present study provides evidence to support the assumption that a disturbance in the homeostasis of electrolytes may also be responsible for difficulties during pregnancy. One possible reaction are premature contractions, for which Fehlinger et al. (1984) showed a correlation to imbalanced overall Mg-levels in a recently published study.

References

- Baltzer, G., E. Daume: Untersuchungen zum Serum-Magnesium-Spiegel in der Gravidität. Verhandlg. Dtsch. Ges. inn. Med. **82** (1976) 880–882.
- Brockhoff, P., M. I. Kurtenbach, W. Stark, K. H. Schicketanz, V. Friedberg, G. H. Rathgen: Klinisch-chemische und haematologische Parameter im Schwangerschaftsverlauf. Gynäk. Rdsch. **21** (1981) 101–123.
- Cushard, W. G., M. A. Creditor, J. M. Canterbury, E. Reiss: Physiologic hyper-

- parathyroidism in pregnancy. J. Clin. Endo. and Metab. **34** (1972) 767–771.
- Davis, R. H., D. B. Morgan, R. S. Rivlin: The excretion of calcium in the urine and its relation to calcium intake, sex and age. Clinical Science **39** (1970) 1–12.
- Di Bona, G. F.: Effect of hypermagnesemia on renal tubular sodium handling in the rat. Am. J. Physiol. **221** (1971) 53–57.
- Domingos, D., C. de Ulhoa, A. B., M. L. Antunes: Magnesium concentration in the blood serum of normal pregnant women. Obstet. Gyn. **25** (1965) 253–255.
- Evans, R. A., L. Watson: Urinary excretion of magnesium in man. Lancet **1** (1966) 522–523.
- Fehlinger, R., C. Kemnitz, P. Dreissig, M. Egert, K. Seidel: Frühgeburtlichkeit, tetanische Reaktionsbereitschaft und Magnesiummangel. Magnesium-Bulletin **6** (1984) 52–59.
- Heaney, R. P., T. G. Skillman: Calcium metabolism in normal human pregnancy. J. Clin. Endocr. **33** (1971) 661–670.
- Leichsenring, J. M., J. Norris, A. Lamson: Magnesium metabolism in college women. J. Nutrition **45** (1951) 477.
- Massry, S. G.: Renal handling of magnesium in the dog. Am. J. Physiol. **216** (1969) 1460–1467.
- Massry, S. G., J. J. Ahumada, J. W. Coburn, Ch. R. Kleeman: Effect of MgCl₂ infusion on urinary Ca and Na during reduction in their filtered loads. Am. J. Physiol. **219** (1970) 881–885.
- Massry, S. G.: Role of hormonal and non-hormonal factors in the control of renal handling of magnesium. Magnesium-Bulletin **3**, 1a (1981) 277–281.
- Rathgen, G. H., P. Brockerhoff, K. H. Schicketanz, V. Friedberg: Klinisch-chemische und hämatologische Parameter. In: Physiologie der Schwangerschaft. Hrsg. V. Friedberg und G. H. Rathgen. Thieme Verlag, Stuttgart 1980, S. 302.
- Sims, E. A. H., K. E. Krantz: Serial studies of renal function during pregnancy and the puerperium in normal women. J. clin. Invest. **37** (1958) 1764.
- Spätling, L.: Orale Magnesium-Zusatztherapie bei vorzeitiger Wehentätigkeit. Geb. Frauenh. **41** (1981) 101–102.
- Spätling, L., P. A. Kunz, D. J. Vonderschmitt, R. Huch, A. Huch: Zum Magnesiumgehalt der Uterusmuskulatur im III. Trimenon. Arch. Gynec. **235** (1983) 470.
- Spätling, L., P. A. Kunz, R. Huch, A. Huch: Magnesium- und Kalziumgehalt der Uterusmuskulatur in der Spätschwangerschaft. In Vorbereitung.
- Warren, E. C., M. D. Wacker, B. L. Vallee: Pathological abnormalities of magnesium metabolism. Med. clin. North America **44** (1960) 1357–1370.

For the authors: Dr. L. Spätling, Ruhr-Universität-Bochum, Frauenklinik Marienhospital, Hölkeskampring 40, D-4690 Herne 1